

Prevalence and management of child sexual abuse cases presented at Nigerian hospitals: A systematic review

Elizabeth Oluwatoyin Akin-Odanye¹

Affiliations:

¹ Ph.D., Clinical Psychology Unit, Department of Family Medicine, University College Hospital, Ibadan, Nigeria.

Corresponding author:

Dr. Akin-Odanye Elizabeth Oluwatoyin, Clinical Psychology Unit, Department of Family Medicine, University College Hospital, Ibadan, Nigeria. E-mail: toyin_akinodanye@yahoo.co.uk

Abstract

Introduction: Aim of this study was to review all suspected sexual abuse (SA) cases presenting at Nigerian hospitals, in order to examine both the prevalence of child sexual abuses (CSA) cases and the availability of hospital protocols and medical and forensic services for their evaluation, management and follow-up.

Methods: A systematic review of literature including only studies published in English from 2006 to 2016 and cited in the Medline, Scopus and Web of Science databases was carried out.

Results: A total of 16 hospital-based studies were included. Out of the total of 166,985 suspected SA cases presented at hospitals, only 1,553 (0.93%) of them were confirmed. Most of them ($n = 1,164$, 74.95%) occurred in children (< 18 years) and majority of the CSA victims were female ($n = 1,155$, 99.23%). HIV screening ($n = 13$, 81.25%) and vaginal swab ($n = 8$, 50%) were the most common investigations carried out by clinicians, whereas the most common medical treatments offered by the health facilities were emergency contraceptives ($n = 8$, 50%) and HIV post exposure prophylaxis ($n = 7$, 43.75%), with only a quarter of the cases ($n = 4$, 25%) offering any form of psychological counseling to victims and/or their parents. Written protocol for CSA evaluation and management was absent in nearly all of the health facilities.

Conclusions: In Nigeria, healthcare and forensic management of CSA are poor, probably due to late presentation of victims at the hospital and to negligence on the part of health and security personnel. CSA is also possibly under-reported, with serious implications for the child and society. Prevention, proper medical treatment, management and follow-up for victims and their families, as well as making the perpetrators answerable for their crimes are highly recommended measures to tackle this issue.

KEY WORDS: Child abuse, sexual; health services; Nigeria; psychological trauma; public health; sexual assault; violence.

Riassunto

Introduzione: L'obiettivo di questo studio è stato quello di fare una revisione di tutti i casi di sospetto abuso sessuale che si sono presentati negli ospedali nigeriani, per esaminare sia la prevalenza dei casi di abuso sessuale nei bambini che la disponibilità di protocolli ospedalieri e servizi medici e forensi per la loro valutazione, gestione e follow-up.

Metodi: È stata effettuata una revisione sistematica della letteratura che ha incluso soltanto studi pubblicati in inglese dal 2006 al 2016 e citati nelle banche dati Medline, Scopus e Web of Science.

Risultati: Un totale di 16 studi su base ospedaliera sono stati inclusi. Dei 166.985 casi totali di sospetta violenza sessuale che si sono presentati in ospedale, solo 1.553 (0.93%) sono stati confermati. La maggior parte di essi ($n = 1.164$, 74,95%) si è verificata nei bambini (< 18 anni) e la maggior parte dei casi di violenza sessuale nei bambini ha interessato soggetti di sesso femminile ($n = 1.155$, 99,23%). Lo screening per HIV ($n = 13$, 81,25%) ed il tampone vaginale ($n = 8$, 50%) sono stati gli esami diagnostici più comunemente richiesti dai medici, mentre le terapie mediche più comunemente somministrate dalle strutture sanitarie sono state la contraccezione d'emergenza ($n = 8$, 50%) e la profilassi post-esposizione per l'HIV ($n = 7$, 43,75%), con una qualche forma di consulenza psicologica offerta alle vittime e/o ai loro genitori soltanto in un quarto dei casi ($n = 4$, 25%). Un protocollo scritto per la valutazione e la gestione dei casi è risultato assente in quasi tutte le strutture sanitarie.

Conclusioni: In Nigeria, la gestione sanitaria e medico legale dei casi di abuso sessuale nei bambini è scarsa, probabilmente per il ritardo nella presentazione delle vittime in ospedale e per la negligenza da parte del personale sanitario e responsabile della sicurezza. I casi di abuso sessuale nei bambini sono anche forse poco denunciati, con una serie di implicazioni per il bambino e la società. La prevenzione, l'appropriato trattamento medico, la gestione ed il follow-up delle vittime e dei loro familiari, così come assicurare alla giustizia i colpevoli di tali crimini sono misure altamente raccomandate per contrastare questo problema.

TAKE-HOME MESSAGE

In Nigeria prevalence of child sexual abuse among sexually-abused cases presented at hospital is high and, possibly, underestimated; however, it is urgent that policy-makers provide funding and health centers implement hospital protocols for giving appropriate medical and forensic management to victims and their families.

Competing interests - none declared.

Copyright © 2018 Elizabeth Oluwatoyin Akin-Odanye Edizioni FS Publishers

This is an open access article distributed under the Creative Commons Attribution (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. See <http://www.creativecommons.org/licenses/by/4.0/>.

Cite this article as: Akin-Odanye EO. Prevalence and management of child sexual abuse cases presented at Nigerian hospitals: A systematic review. J Health Soc Sci. 2018;3(2):109-124

DOI 10.19204/2018/prv12

Received: 26/03/2018

Accepted: 19/04/2018

Published: 15/07/2018

INTRODUCTION

The Child Rights Act adopted by Nigeria in 2003 [1] requires that children's well-being must be respected and considered paramount. However, the increased incidence of child sexual abuse cases has dominated the headlines in the media, causing a great deal of concern. Child sexual abuse (CSA) describes all acts directed at engaging a child directly or indirectly in non-age appropriate sexual behaviours either through intimidation or inducement. The World Health Organization (WHO) has defined child sexual abuse as 'the involvement of a child in sexual activity that he or she does not fully comprehend, is unable to give informed consent to, or for which the child is not developmentally prepared and cannot give consent, or that violates the laws or social taboos of the society' [2]. Minahan considers CSA as occurring at an age when the child does not fully understand the nature of the act taking place and, therefore, lacks the cognitive, emotional and physical power to say no to the sexual behavior [3]. Features that characterize CSA include but are not limited to: asking or pressuring a child to engage in sexual activities (whether successful or not); indecent exposure of the genitals to a child; and child exposure to or involvement in pornography, physical (fondling, caressing, penetration, etc) or non-physical sexual contact with the child's genitals or other body parts.

The occurrence of CSA is a global and widespread phenomenon. A literature review showed that CSA is the third most common type of child maltreatment [4]. The 2009 WHO's report showed an annual global estimate of CSA at about 73 million (7%) for boys and 150 million (14%) for girls, affecting 1 in 4 girls and 1 in 6 boys before the age of 18 [5]. In 2008, the Centers for Disease Control and Prevention (CDC) reported that 772,000 children were abused or neglected all over the world, and approximately 9% of them were sexually abused [6]. The United States Department of Health and Human Services (USDHHS) reported that, in 2010, over 60,000 American children were sexually

abused [7].

Africa is not an exception and, according to a WHO report, sexual abuse of females before the age of 15 accounts for between 7% and 21% of all CSA cases, whereas figures for adolescent males ranged between 3.4%-29.9% in some African countries [8]. Also, more than 41% of rapes reported in South Africa involved children under age 18 and it was suggested that 25% of girls are likely to be raped before the age of 16 [9]. In one Ugandan study, 72% of victims who presented to a hospital were aged 12 or below [10]. Similarly, a Kenyan study found that 50% of patients presenting to some sexual assault centres were under 14 years old [11]. Gwiryayi reported an overall CSA prevalence rate of 56.3% among children attending urban secondary schools in Zimbabwe [12]. Likewise, an analysis of Demographic and Health Survey's data on 6,351 adolescents aged 18 years or below from six countries in sub-Saharan Africa, between 2006 and 2008, showed that prevalence of CSA ranged from 1.04% in Liberia to 5.84% in Zambia, with Uganda, Nigeria, Ghana, and Zimbabwe reporting a prevalence of 1.38%, 2.40%, 4.61%, and 4.96%, respectively [13].

Community-based surveys among in-school and out-of-school adolescents in Nigeria indicated high levels of CSA. In a study from Maiduguri, North-East Nigeria, a sexual assault rate of 77.7% was reported among female children workers with sexual assault being more likely in girls who were younger than 12 years [14]. Also, Kunuji and Essiet reported that about 14% and 35% of out-of-school adolescents in an urban slum in Lagos had been victims of rape and statutory rape, respectively [15]. Olley [16] and Manyike et al. [17], reported a prevalence of 55% and 40% among in-school adolescents in Southwest and South-east Nigeria, respectively. Other studies were based on newspaper accounts indicating a high incidence of CSA in different parts of Nigeria [18, 19]. In Benin City, victims of sexual molestation comprised children aged 6-12 years and adolescents aged 13-19 years [20]. Another study showed

that 58% of rape cases reported in hospitals in Benin City concerned child victims [21]. Although female children were reported in literature as predominantly being sexually abused, incidence of sexual abuse on male children is on the increase as well [22].

CSA can take several forms. Giving away of young girls into marriage even before the onset of their menstrual cycle is still being practiced especially in the Northern part of Nigeria for religious and economic reasons. Child sexual abuse can manifest itself very frequently in the form of child prostitution. In this way, Uzuegbu observed the existence of a growing market for the trafficking of women and young children from Nigeria to Europe, America and some African countries for the illicit purposes of prostitution and pornography [23]. Female genital mutilation (FGM) referring to all procedures that involve partial or total removal of the external female genitalia, or other injury to the female genital organs for non-medical reasons, is another form of CSA that is rampant in Nigeria. The news media are getting flooded with cases of intra-familial child sexual abuses especially in the father-daughter relationships. Indeed, the majority of children and teen victims know the perpetrators, who are mostly male and often relatives, neighbours, or other trusted members of their social network [24, 25].

Low levels of disclosure in CSA victims have been also reported in literature [26–28]. Reasons for poor disclosure range from shame, anxiety, or fear [29] to likelihood of stigma, ridicule and/or reduction of their chances of getting married if the abuse was made public, especially amongst the female victims. The first point of contact for victims of sexual assault who decide to seek redress for the sexual infringement may be a police station or a medical facility, such as an emergency centre or an outpatient health centre. The type of reception and quality of service received by those who were bold enough to report their ordeal to law enforcement agents or visit healthcare facilities for treatment, will most likely determine if victims will complete follow-up visits as recommended or if others with whom they

share their experience will be willing to use or recommend the same available services. This research, therefore, aimed to systematically review retrospective studies focusing on child sexual abuse cases presented at Nigerian hospitals, with the specific objectives of determining the prevalence of this phenomenon, the availability of hospital protocols for the evaluation, management and follow-up of CSA cases as well as the availability of medical and forensic services to CSA victims. The findings could give more valuable information to policy makers, in order to improve short and long-term measures for both clinical and forensic assistance to CSA victims and their families, in Nigeria as well as other sub-Saharan countries.

METHODS

In order to carry out this review, we followed the statements of Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) [30].

Study selection

Articles in scientific journals, excluding doctoral theses or book chapters, published only in English, and focused on studies reporting hospital-based cases of sexual abuse including CSA, presented for treatment at Nigerian hospitals, were retrieved using the Medline, Scopus, and Web of Science databases. Abstracts of all suitable articles were examined. The same selection criteria were applied for checking references of the selected publications to identify further reports not found by database searching. The search terms including '*sexual abuse*' OR '*sexual assault*', OR '*child rape*' AND '*child*', and '*management*' AND '*child sexual abuse*', and '*hospital-based*' AND '*child sexual abuse*' with the term '*Nigeria*', were followed. Only studies published from 2006 to 2016 were considered.

Inclusion and exclusion criteria

In order to be included in this review, studies had to meet the following criteria: (A) focus on a sample of children and adolescents, including studies in which the mean age of par-




Figure 1. Flowchart of the selection of studies for inclusion in the review.

ticipants was less than 18 years, or the upper limit of the age range of the sample was less than 41 years, if the study did not report the mean age of participants; (B) provide prevalence data on hospital-based childhood sexual abuse cases in Nigeria; (C) with regard to the abuses, provide data on types of hospital protocols and treatment provided for the evaluation, management and follow-up of CSA cases; d) provide details about investigations, and/or legal actions taken against perpetrators. Studies that did not meet the inclusion criteria were excluded from this review.

Study coding

A series of variables relating to the studies reviewed were recorded: (a) author(s) and year of publication; (b) study design; (c) Nigeria's geopolitical zone. In addition, content was analyzed in order to identify: 1) Prevalence of SA cases among adults and children reported by retrieved hospital-based studies; 2) Investigations carried out and treatment provided

to the victims; 3) Involvement of law enforcement agencies in prosecuting offenders. Each study was analyzed by the absolute (n) and relative (%) frequency for variables as follows: 1) Number of all suspected SA cases presented at hospital; 3) Prevalence of SA cases confirmed in adults and children (CSA); 4) Prevalence of CSA grouped by gender; 4) Mean age of confirmed cases of CSA; 5) Presence or absence of: i) Hospital protocol for evaluation, management and follow up of CSA cases; ii) Psycho-social support for CSA victims and/or their families; iii) Screening for Sexually Transmitted Diseases (STDs) and HIV; iv) Post-exposure prophylaxis for STDs and HIV; v) Pregnancy test and emergency contraception.

RESULTS

The process of selecting the studies in this review is shown in Figure 1. After the initial search (101 potential articles), 16 observational, retrospective studies were included;

Table 1. Authors, period covered in review, geopolitical zones, states, hospitals and departments represented in reviewed studies ($n = 16$ studies).

Authors and period covered	Type of study design	Geopolitical zones, state, city, hospitals and department
Abdulkadir et al., 2011 (1 st Jan 2007 - 30 th June 2008) ³¹	Cross-sectional	<i>North Central Nigeria</i> GOP General Hospital, Suleja, Niger State.
Daru et al., 2011 (Jan 2001 - Dec 2003) ³²	Cross-sectional	Gynaecological Emergency Unit JUTH, Jos, Palteau State.
Ashimi et al., 2015 (Sept 2011 - Aug 2013) ³³	Longitudinal	<i>North West Nigeria</i> Gynecological Emergency Unit of a Tertiary Health Facility in a Birnin-Kudu, Jigawa State.
Bugaje, et al., 2012 (Aug 2008 - Oct 2009) ³⁴	Cross-sectional	ICH, Zaria, Kaduna State.
Hassan et al., 2016 (Jan 2010 - Dec 2014) ³⁵	Cross-sectional	UDTH, Sokoto, Sokoto State.
Chinawa et al., 2013 (Jan - Dec 2010) ³⁶	Cross-sectional	<i>South East Nigeria</i> CHOP EBSUTH Abakiliki, Ebonyi State.
Ekabua et al., 2006 (Jan 1998 to Dec 2001) ³⁷	Cross-sectional	Department of Obstetrics and Gynaecology, University of Calabar Teaching Hospital, Calabar, Cross River State.
Nwolisa et al., 2016 (Jan 2009 to Dec 2013) ³⁸	Cross-sectional	Paediatric Infectious diseases unit, Federal Medical Centre, Owerri, Imo State, Nigeria.
Olatunya et al., 2013 (Jan 2010 - March 2013) ³⁹	Cross-sectional	<i>South West Nigeria</i> CHOP, EKSUTH, Ado-Ekiti, Ekiti State.
Akinlusi et al., 2014 (Jan 2008 - Dec 2012) ⁴⁰	Cross-sectional	Gynaecology Clinic LASUTH, Ikeja. Lagos State.
Badejoko et al., 2014 (1 st Jan 2007 - 31 st Dec 2011) ⁴¹	Cross-sectional	Children/Adult Emergency units OAUTHC, Ile-Ife, Osun State.
Ige & Fawole, 2012 (June 2008 - May 2009) ⁴²	Cross-sectional	General Hospital in Ibadan, Oyo State.
Adeleke et al., 2012 (1 st Jan 2003 - 31 st Dec 2009) ⁴³	Cross-sectional	State hospital, Asubiaro, Osogbo, Osun State.
Ezechi et al., 2016 (Jan 2006 - Dec 2015) ⁴⁴	Cross-sectional	HIV treatment centre, Nigerian Institute of Medical Research, Lagos.
Duru, et al., 2014 (1 st Dec 2009 - 30 th Nov 2013) ⁴⁵	Cross-sectional	<i>South South Nigeria</i> CHOP, NDUTH, Okolobiri, Bayelsa State.
Akhiwu et al., 2013 (1 st Jan 2000 - 31 st Dec 2000) ⁴⁶	Cross-sectional	Police Medical Centre, Benin City, Edo State.

Notes: GOP – General Outpatient Dept.; CHOP – children out patient / pediatric outpatient; EBSUTH – Ebonyi State University Teaching Hospital; UDTH - Usman Danfodio Teaching Hospital; (OAUTHC - Obafemi Awolowo University Teaching Hospitals complex; LASUTH - Lagos State University Teaching; NDUTH – Niger Delta University Teaching Hospital Okolobiri; EKSUTH - Ekiti State University Teaching Hospital; JUTH - Jos University Teaching Hospital, ICH – Institute of Child Health.

Table 2. Prevalence of child sexual abuse cases reported in Nigerian health institutions ($n = 16$ studies).

Authors' references	All cases presented at hospitals		Total of CSA cases (< 18 years)			Prevalence of confirmed CSA cases ($n =, %$)
	All cases (n)	All confirmed SA cases ($n =, %$)	Boys (n)	Girls (n)	Mean age in years	
Daru et al., 2011 ³²	2,135	105 (4.9%)	0	67	12	67 (63.8%)
Abdulkadir et al., 2011 ³¹	No data Available	81 (ID)	0	77	9	77 (95.06%)
Ige and Fawole, 2012 ⁴²	No data Available	72 (ID)	0	72	-	72 (100%)
Bugaje et al., 2012 ³⁴	33,313	20 (0.06%)	3	17	-	20 (0.06%)
Akhiwu et al., 2013 ⁴⁶	1,028	85 (4.67%)	0	48	14.54	48 (56.47%)
Chinawa et al., 2013 ³⁶	3,750	33 (0.9%)	2	31	-	33 (0.9%)
Olatunya et al., 2013 ³⁹	6,535	28 (0.43%)	0	28	-	28 (0.43%)
Duru et al., 2014 ⁴⁵	12,229	33 (0.27%)	1	32	7.9	33 (0.27%)
Akinlusi et al., 2014 ⁴⁰	39,770	287 (0.6%)	0	240	12.9	240 (83.62%)
Badejoko et al., 2014 ⁴¹	24,575	76 (0.13%)	0	31	-	31 (40.79%)
Ashimi et al., 2015 ³³	973	24 (2.26%)	0	22	8.8	22 (91.67%)
Hassan et al., 2016 ³⁵	5,317	34 (0.39%)	0	21	12.6	21 (61.76%)
Adeleke et al., 2012 ⁴³	14,970	309 (2.06%)	0	228	15.8	228 (73.79%)
Ekabua et al., 2006 ³⁷	1,038	22 (2.11%)	0	15*	13.6	15 (68.18%)
Ezechi et al., 2016 ⁴⁴	21,199	196 (0.9%)	2	79**	-	81 (41.33%)
Nwolisa et al., 2016 ³⁸	153	148 (96.7%)	1	147	13.33	148 (96.7%)
Total	166,985	1,553 (0.93%)	9	1,155	-	1,164 (74.95%)***

Notes: *SA* – Sexual Abuse; *CSA* – Child Sexual Abuse; * 15 victims \leq 14 years; ** 81 victims < 20 yrs, since age of the 12 male victims ranged from 19 to 41 years with a mean of 24.9 ± 4.9 years⁴⁴ it was assumed that 2 males were 19 years; *** All SA cases/Total CSA cases presented in hospitals; ID = Insufficient Data.

the articles that did not meet the previously established inclusion and exclusion criteria were excluded.

Majority of the selected articles were cross-sectional and retrospective studies, except one that was longitudinal (prospective) study. The papers reviewed were from 5 out of the 6 geopolitical zones in Nigeria. No study was found during the period of the review from the North East geopolitical zone. More specifically, 6 studies were from hospitals in South-West Nigeria (Ile-Ife, Ibadan, Ado-Ekiti, Oshogbo and Lagos-2 studies), 3 from North-West (Jigawa, Kaduna, Sokoto), 2 each from North Central (Niger, Plateau) and South-South (Bayelsa, Edo), and 3 from South-East (Ebonyi, Calabar and Owerri) Nigeria (see Table 1).

With regard to the 36 states of Nigeria plus the Federal Capital Territory (FCT), 16 studies were reviewed from 14 states covering 37.8% of the Nigerian territory, including

the FCT. The papers reviewed were published between 2006 and 2016, but years covered by studies reviewed ranged from 1998 to 2015.

Prevalence of child sexual abuse cases presented at hospitals

The prevalence of all sexual abuse cases among patients presented at hospitals ranged from 0.06% in Zaria [34] to 96.7% in Owerri [38]. A summary of all sexual abuse cases and percentage of CSA with respect of all sexual abuse cases drawn from the 16 hospital-based studies included in this review are shown in Table 2.

As shown in Table 2, out of the estimated total of 166,985 suspected cases of sexual abuse that presented at hospitals, only 1,553 (0.93%) of them were confirmed as being sexual abuse cases. Moreover, of the 1,553 sexual abuse cases reported, 1,164 (74.95%) were CSA as they occurred in children aged 18 years or below. Majority ($n = 1,155, 99.2\%$) of the




Figure 2. Types of medical investigations carried out in reviewed studies ($n = 16$).

sexually abused children were female. This displays a high vulnerability of the Nigerian girl child to sexual abuse. It could, however, mean that most cases involving the female children presented to the hospital for treatment, because the physical damage which may have been caused on genitals is often more obvious and severe than compared to what may obtain in the male child (except in cases of anal penetration). Similar findings about the prevalence of child sexual abuse cases presented at Nigerian hospitals have been reported in studies from other Sub-Saharan countries. For instance, an Ethiopian hospital-based study on 275 sexually and physically abused children, confirmed that majority of cases were victims of sexual abuse (97.3%) [47]. Moreover, in a hospital-based prospective study from Uganda, about 92% of sexual assault cases were against children aged 1-17 years [48]. Foumane et al., (2014) [49] also reported a high prevalence (65%) of CSA in Yaoundé, Cameroon. Another study from Dammam, Saudi Arabia, showed that 85% of the 87 cases of assaults on children aged less than 18 at the Maternity and Children Hospital (from 2008 to 2010), and at Forensic and Legal Center (from 2006 to 2010) were sexually motivated [50]. A much lower incidence of 15 children aged 6-12 years was reported in a study conducted by the Forensic Medical Au-

thority (FMA) of the Fayoum Governorate in Egypt over a 5-year period [51]. However, unlike the findings highlighted in this Nigerian hospital-based review, 80% of the victims of child sexual assault in Egypt were males and only 20% were females [51]. Other studies from Ethiopia [47, 52, 53] and Malawi [54] have reported findings similar to those discovered by this systematic review, as a higher proportion of female than male children were victims of child sexual abuse.

Management of sexual abuse cases in Nigerian hospitals

The most commonly requested investigations across the 16 studies were as follows: a) HIV screening ($n = 13$, 81.25%); b) Vaginal swab ($n = 8$, 50%); c) HBsAg ($n = 6$, 37.5%); d) Hepatitis C ($n = 4$, 25%); e) Pregnancy test ($n = 5$, 31.3%); f) Tetanus toxoid, urine culture, abdominal ultrasound and VDRL ($n = 2$, 12.5%); g) Full blood count and PCV ($n = 1$, 6.3%).-

Data from the current review showed that important investigations for sexual abuse cases presented at Nigerian hospitals are not routinely requested for across all Nigerian hospitals that were analyzed in this review. Moreover, even among those studies where investigations were requested, only few victims actually carried out the requested in-

Table 3. Available services and procedures in the management of child sexual abuse ($n = 16$ studies).

Authors' references	CSA Hospital protocol	Psychosocial support services	Screening		PEP		Pregnancy		Follow-up
			STDs	HIV	STDs	HIV	Test	EC	
Daru et al., 2011 ³²	Not available	Not offered	Yes 61%	Yes 64%	NDS	NDS	NDS	Yes 23%	NDS
Abdulkadir et al., 2011 ³¹	Not available	NDS	NDS	NDS	NDS	NDS	NDS	NDS	NDS
Ige and Fawole, 2012 ⁴²	NDS	Offered to 15.3% victims	Yes 76%	Yes 60%	Yes 47%	Yes 6%	ND	Yes 12%	No 0%
Bugaje, et al., 2012 ³⁴	NDS	Not offered	Yes 65%	Yes 95%	Yes 50%	NDS	Yes 15%	NDS	Yes 45%
Akhiwu et al., 2013 ⁴⁶	NDS	NDS	NDS	NDS	NDS	NDS	NDS	NDS	NDS
Chinawa et al., 2013 ³⁶	NDS	Not offered	Yes 30%	Yes 42%	NDS	NDS	No	No	Yes 3%
Olatunya et al., 2013 ³⁹	Not available	NDS	NDS	Yes 100%	Yes 89%	Yes 61%	No	Yes 44%	Yes 3.6%
Duru, et al., 2014 ⁴⁵	NDS	Not offered	Yes 27%	Yes 62%	NDS	Yes 33%	NDS	Yes 12%	Yes 18.2%
Akinlusi et al., 2014 ⁴⁰	Not available	Not offered	Yes 64%	Yes 74%	NDS	Yes 29%	Yes 31%	Yes 22%	Yes 72%
Badejoko et al., 2014 ⁴¹	Not available	NDS	Yes 42%	Yes 85%	Yes 92%	Yes 52%	Yes 13%	Yes 37%	Yes 13%
Ashimi et al., 2015 ³³	NDS	Offered Pre STD/HIV screening counselling	Yes 13%	Yes 46%	NDS	NDS	NDS	NDS	0% after 2 nd visit
Hassan et al., 2016 ³⁵	NDS	NDS	NDS	NDS	NDS	NDS	NDS	NDS	NDS
Adeleke et al., 2012 ⁴³	Not available	NDS	Yes 2%	Yes 2%	No 0%	No 0%	Yes 0.6%	NDS	Yes 3.9%
Ekabua et al., 2006 ³⁷	NDS	Not offered	Yes 14%	Yes 27%	NDS	NDS	Yes	Yes 27%	No
*Ezechi et al., 2016 ⁴⁴	Available	Offered	Yes 34%	Yes 100%	NDS	Yes 90%	NDS	Yes 95%	Yes 75%
Nwolisa et al., 2016 ³⁸	NDS	Offered	NDS	Yes 100%	NDS	Yes 90%	NDS	NDS	Yes 1.4%

Notes: EC = Emergency contraception; NDS = Not documented in the study; PEP = Post Exposure Prophylaxis; *All cases of occupational and non-occupational exposure to infectious agents are provided services using a standardized management protocol and 75% completed 3months follow-up visits.⁴⁴

vestigations. For instance, in one of the reviewed studies, 29 HIV screening test were requested, but only 18 of them were performed and of 20 vaginal swabs requested only 5 were actually carried out [45]. Reasons for this non-compliance might be due to lack of financial resources available to the victims and their parents.

As reported by the 16 hospital-based studies reviewed, the most common treatments offered to children who were victims of sexual abuse included: emergency contraceptives

($n = 8$, 50%), HIV post-exposure prophylaxis (PEP) ($n = 7$, 43.75%), antibiotics ($n = 4$, 25%), and some types of psychosocial support ($n = 4$, 25%). It is noteworthy that 56% and 75% of the studies reviewed did not document, nor did routinely offer HIV and sexually transmitted diseases (STDs) prophylaxis, respectively. A study in Uganda also reported that HIV prophylaxis was not given to sexual abuse victims due to the non-availability of funds [48]. The low proportion of children who received PEP for HIV and other STDs

was also likely due to delayed arrival of the victims at hospital. In most of the studies reviewed, a good number of the CSA victims went to hospital 72 hours after the occurrence of the sexual abuse [36, 42, 45]. However, in two of the studies reviewed, a high proportion of CSA cases presented at health centers within 72 hours and 24 hours, respectively [35, 41]. Prompt presentation, within 72 hours, is essential for proper management of sexual abuse or assault cases. Delayed presentation might be, however, attributed to low awareness of ideal timing for STDs and HIV prophylaxis, for emergency contraceptives, as well as due to financial constraints and geographic barriers [55]. Indeed, probably prophylaxis is not often available at health centers, which are dedicated to the provision of HIV and STD prophylaxis to high-risk patients due to lack of funds. According to Patel et al., some elements of a comprehensive medical care management of sexual abuse cases include: 1) Immediate attention to acute physical trauma; 2) History and physical examination; 3) Immediate and follow-up mental health counseling; 4) Tests and prophylaxis for sexually transmitted infections and HIV; 5) Counselling and provision of emergency contraception [56]. The management of sexual assault is based on the principles of preventing pregnancy, treating CSA complications and providing prophylactic antibiotics, as well as psychological rehabilitation [32]. Unfortunately, these principles were not strictly followed by the hospital-based studies reported by this review. Section 32, subsections 1 and 2 of the Nigerian Child Right Act apart from providing a penalty of 14 years imprisonment for persons who sexually abuse or sexually exploits children in any manner, requires mandatory counseling and rehabilitation of abused children [1]. However, in most of the studies reviewed, the CSA victims were not referred for psychological support [34, 40, 45]. Only few studies reported that some form of counseling was provided to the CSA victims, by receiving a specific treatment at health facility [33, 42, 38, 44]. Among the 138 resident physicians in gynecology and obstetrics

attending an update course in Nigeria, only 39 (28.7%) of them indicated access to clinical psychologists in their health facilities [57]. It is not clear how much of the available clinical psychologists are involved in the care of rape and child sexual abuse victims. Sexual violence has short- and/or long- physical and psychological negative effects on its victims. Psychologically, victims of CSA are more likely to experience depression, substance abuse, increased risk for adoption of risky sexual behaviours, post-traumatic stress disorder (PTSD), and suicide [16, 58]. Physically, multiple bruises in uncommon sites, vaginal and anal lacerations and perforations as well as exposure of victims to unwanted pregnancy and sexually transmitted infections (STIs) including HIV/AIDS have been reported [36, 45]. As shown in literature, over their lifetime, sexual assault survivors are at increased risk for post-traumatic stress disorder (30%), major depression (30%), suicidal ideation (33%), and suicidal attempt (13%) [58]. Psychological interventions should be incorporated into healthcare services provided to CSA victims. Thought must also be given to providing support and/or counselling to those caring for the child. This may be required even if the CSA victim is not assessed as needing therapy. Early interventions that help individuals through their initial reactions to sexual assault could reduce or prevent more severe psychological distress. Therefore, immediate trauma management and long-term counseling and psychotherapy are essential components of CSA survivor care in moderating the negative consequences of CSA on victims.

Presence of hospital protocols for CSA cases

With regard to the hospital protocols, this review found that written protocols for CSA evaluation, management and follow-up were not available in most of the medical facilities [31, 34, 40]. This finding is in line with that reported by a study on 138 resident physicians in gynecology and obstetrics across 25 different training institutions in Nigeria, from all 6 geopolitical zones, where 131 of

them (94.9%) claimed a lack of rape management protocol in their training health facilities [57]. The less than desirable standard of medical management of the CSA victims could be linked to this lack of protocol that could serve as a ready reference point on what has to be done in case of CSA. Furthermore, most of the hospitals were poorly equipped to adequately manage CSA cases, probably due to inadequate funding by government.

Overall, most of the studies reviewed have reported poor follow-up practices on CSA cases. Parents need to be educated from the initial visit on the importance of follow-up after the initial contact with the health facility to ensure proper investigations and treatment that could prevent unwanted health

outcomes. Follow-up visits could also create an avenue to provide psychosocial care in the form of ascertaining adequate support for the child and family, as well as an opportunity to provide child sexual abuse prevention and safety teaching to children and families.

Prosecution of child sexual abuse cases in Nigeria

In this review, although a good number of assailants were known, only few arrests were made. In a study where 49 (69.0%) victims reported their ordeal to the police, only 24 (33.8%) perpetrators were arrested [41]. In another study, only 5 (21.7%) out of 23 perpetrators caught were prosecuted [45]. In

Table 4. Prosecution of Child Sexual Abuse cases reported to the police ($n = 16$ studies).

Authors' references	All SA cases (n)	Assailants known (n, %)	Forensic specimen collected (n, %)	Cases reported to police (n, %)	Assailants arrested (n, %)	Prosecutions (n, %)	Assailants sentenced (n, %)
Daru et al., 2011 ³²	105	81 (77.4%)	NDS	NDS	NDS	NDS	NDS
Abdulkadir et al., 2011 ³¹	81	79 (97.5%)	NDS	NDS	NDS	NDS	NDS
Ige & Fawole, 2012 ⁴²	72	45 (62.5%)	NDS	17 (23.6%)	NDS	NDS	NDS
Bugaje, et al., 2012 ³⁴	20	14 (70%)	NDS	12 (60%)	NDS	NDS	NDS
Akhiwu et al., 2013 ⁴⁶	85	75 (88.2%)	NDS	ALL	NDS	6 (7.1%)	NDS
Chinawa et al., 2013 ³⁶	33	NDS	NDS	6 (18.1%)	NDS	NDS	NDS
Olatunya et al., 2013 ³⁹	28	NDS	NDS	17 (60.7%)	NDS	0	NDS
Duru, et al., 2014 ⁴⁵	33	23 (67.7%)	NDS	5 (21.7%)	NDS	NDS	NDS
Akinlusi et al., 2014 ⁴⁰	287	210 (73.1%)	No	265 (92.2%)	NDS	NDS	NDS
Badejoko et al., 2014 ⁴¹	76	47 (62%)	No	52 (69%)	25 (33.8%)	NDS	NDS
Ashimi et al., 2015 ³³	24	20 (83.3%)	NDS	NDS	NDS	NDS	NDS
Hassan et al., 2016 ³⁵	34	20 (58.8%)	NDS	20 (58.8%)	14 (41.2%)	NDS	NDS
Adeleke et al., 2012 ⁴³	309	246 (79.6%)	NDS	ALL	NDS	NDS	NDS
Ekabua et al., 2006 ³⁷	15	NDS	NDS	NDS	NDS	NDS	NDS
Ezechi et al., 2016 ⁴⁴	81	42 (52%)	NDS	NDS	NDS	NDS	NDS
Nwolisa et al., 2016 ³⁸	148	61 (41.2%)	NDS	NDS	NDS	NDS	NDS

Note: NDS = Not documented in the study.

some other studies, where there were police involvements no prosecution was made [34, 39]. In Nigeria, literature statistics reported few cases of child sexual abuse, probably due to the social stigma that such a report would earn the victim. However, in Nigeria there are not just few police reports, there are also few trials of perpetrators of child sexual abuse. Instances of inadequate prosecution of CSA cases in Nigeria including overtly lenient sentence granted CSA abuse perpetrators by our judicial system abound [19]. For instance, the Lagos state government with a population of almost 20 million reported 283 cases of child defilement in 2011, out of which only 10 were prosecuted and convicted, which is less than a 10% conviction rate [59].

Some reasons for the underreporting and poor prosecution of CSA perpetrators is associated with the culture of blaming the victim and the desire of victims and/or their families to avoid shame and stigma they often opt to settle out of court with the consequence. As noted by Duru et al. [45], most cases of CSA, especially when perpetuated within the family, remain undisclosed in many parts of Africa. Also, government-based care and law enforcement agencies that ought to support the CSA victims and ensure they get justice are often lacking in skills and commitment. Police officers often lack the requisite specialized training that enable them to handle CSA cases in empathetic and compassionate manner, as well as skills needed to successfully carry out forensic identification of alleged perpetrators. Legislatively, the penalty for acts of CSA is often not severe enough to deter perpetrators.

Limitations of the Study

A major limitation of this study is due to incomplete data existing in most of the retrospective studies reviewed. Also, age of respondents was not routinely categorized in a way that number of CSA victims may be demarcated from adults. Despite these limitations, the findings reported in this review could give a good representation of the state of clinical and forensic services provided to child sexual

abuse victims in Nigerian hospitals.

Recommendations and take-home messages
There is a high prevalence of CSA cases among sexually-assaulted cases presented at Nigerian hospitals.

Child sexual abuse victims and their families receive suboptimal medical care due to lack of protocols for the evaluation and management of child sexual abuse cases, and due to poor funding of healthcare facilities.

Currently, only few cases of child sexual abuse get reported to the police and even fewer get to court due to poor forensic skills and social stigma.

There is an urgent need to educate Nigerian people about the importance of presenting to hospitals immediately after the occurrence of violence, for receiving appropriate medical treatment and promoting the needed collection of viable forensic evidence that could facilitate the prosecution of sexual abuse perpetrators.

CONCLUSION

There is a high prevalence of CSA cases among reported cases of sexual violence presented at Nigerian hospitals. Social, health, forensic, legislative and judicial services that can deter this undesirable act and facilitate the effective rehabilitation of victims are grossly inadequate. Government agencies, parents, and all responsible adults should help protect children from perpetrators through adequate education and conscious effort to avoid putting children in vulnerable situations. In agreement with Badejoko et al., most acts of CSA in Nigeria are perpetrated by well-known and trusted familiar called for caution by parents and guardians when leaving children alone. Therefore, it is urgent that both the law enforcement agencies and the legislative and judicial arms of government wake up to their responsibility and work cooperatively for ensuring that CSA perpetrators get penalties that are commensurate with this type of crime. Most of the studies reviewed reported the lack of CSA care protocols in Nigeria and inadequate funding resulting in ill-equipped hospitals as the reason for lapses in the hospi-

tal care of CSA victims. It is essential that Nigerian hospitals develop a protocol to aid the efficient provision of care for victims of sexual abuse. Such a protocol should be developed by multidisciplinary teams to promote acceptability and allow for predictability of flow of the sexually-abused patients from one health care centre to another within the healthcare system. Necessary training of healthcare providers to effectively implement this protocol would also be required. The success of SA treatment protocols used in Kenya shows that creating and using similar models is feasible also in other sub-Saharan African states [11]. In situations where the government is unre-

sponsive to the needs of hospitals in terms of funding, hospital administrators and pediatric departments should be proactive in sourcing funds through public private partnerships and applying for grants from children-focused local and international agencies. In conclusion, prevalence of child sexual abuse among sexually-abused cases presented at Nigerian hospital is high and, possibly, underestimated; however, it is urgent that policy-makers provide funding and policy makers implements hospital protocols for giving appropriate medical and forensic management to victims and their families.

References

1. UNICEF. Child Right Act (2003) [cited 2018 Dec 10]. Available from: https://www.unicef.org/nigeria/ng_publications_Childs_Right_Act_2003.pdf.
2. World Health Organization. Violence & Health in the WHO African Region. Brazzaville: World Health Organization Regional Office for Africa; 2010 [cited 2018 Dec 10]. Available from: <http://www.afro.who.int/sites/default/files/2017-06/mvi-violence-health-15-04-11.pdf>.
3. Minahan A. Encyclopaedia of Social Work. Vol. 1 Mary land: NASW; 1987.
4. Law Enforcement. In the 2011 statistical abstract: The national data book; 2011 [cited 2018 Dec 10]. Available from: <http://www.census.gov/compendia/statab/2011/tables/11s0339.pdf>.
5. World Health Organization. Prevention of child maltreatment; Geneva:WHO;2009 [cited 2018 Dec 10]. Available from: www.who.int/bulletin/volumes/87/5/08-057075.pdf.
6. National Center for Injury Prevention and Control (U.S.). Division of Violence Prevention. Child maltreatment: facts at a glance (2010). National Center for Injury Prevention and Control (U.S.). Division of Violence Prevention; 2010 [cited 2018 Dec 10]. Available from: http://www.cdc.gov/ViolencePrevention/pub/CM_datasheet.html.
7. US Department of Health and Human Services. (2012). Testimony [cited 2018 Dec 10]. Available from: http://www.hhs.gov/asl/testify/2011/12/t20111213a.html#_ftnref1.
8. WHO. WHO multi-country study on women's health and domestic violence against women. Geneva: World Health Organization; 2005 [cited 2018 Dec 10]. Available from: whqlibdoc.who.int/publications/2005/9241593512_eng.pdf.
9. Cox S, Andrade G, Lungelow D, Schloetelburg W, Rode H. The child rape epidemic: assessing the incidence at Red Cross Hospital, Cape Town, and establishing the need for a new national protocol. *South Afr Med J.* 2007;97:950–955.
10. Ononge S, Wandabwa J, Kiondo P, Busingye R. Clinical presentation and management of alleged sexually assaulted females at Mulago hospital, Kampala, Uganda. *Afr Health Sci.* 2005;5:50–54.
11. Ranney ML, Rennert-May E, Spitzer R, Chitai MA, Mamlin SE, Mabeya H. A novel ED based sexual assault centre in Western Kenya: description of patients and analysis of treatment patterns. *Emerg Med J.* 2011;28:927–931.
12. Gwirayi P. Child Sexual Abuse among Urban Secondary School Pupils: Impact of Family Characteristics and Family Structure. *Int Rev Soc Sci Humanit.* 2012;3(1):36–50.

13. Yahaya I, Uthman OA, Soares J, Macassa G. Social disorganization and history of child sexual abuse against girls in sub-Saharan Africa: a multilevel analysis. *BMC Int Health Hum Rights*. 2013;13:33.
14. Audu B, Geidam A, Jarma H. Child labor and sexual assault among girls in Maiduguri, Nigeria. *Int J Gynecol Obstet*. 2009;104:64–67.
15. *Kunnuji MO, Esiet A*. Prevalence and Correlates of Sexual abuse among female Out-of School Adolescents in Iwaya Community, Lagos State, Nigeria. *Afr J Reprod Health*. 2015;19(1):82–90.
16. Olley BO. Child sexual abuse as a risk factor for sexual risk behaviours among socially disadvantaged adolescents in Ibadan, Nigeria. *Vulnerable Children and Youth Studies*. 2008; 3(3):243–248.
17. Manyike PC, Chinawa JM, Aniwada E, Odutola OI, Chinawa TA. Child sexual abuse among adolescents in southeast Nigeria: A concealed public health behavioral issue. *Pak J Med Sci*. 2015;31(4):827–832. doi:10.12669/pjms.314.7115.
18. Okonkwo O, Naish ME. *Criminal Law in Nigeria*. Ibadan: Spectrum Publisher;2013.
19. Akpoghome TU, Nwano TC. Examining the Incidences of Sexual Defilement of Children in Nigeria Donnish. *J Law Conflict Resol*.2016; 2(1):1–9.
20. Olusanya O, Ogbemi S, Unuigbo J, Oronsaye A. The pattern of rape in Benin city, Nigeria. *Trop Geog Med*. 1986;38(3):215–220.
21. Omorodion FI, Olusanya O. The social context of reported rape in Benin city, Nigeria. *Afr J Reprod Health*.1998;2(2):37–43.
22. Bejide F. Male Child Victim of Sexual Abuse in Nigeria. *Front Leg Res*. 2014;2(1):83–99.
23. Uzuegbu J. A Critical Survey of Some Societal Violence Against Women and Young Girls: The Nigerian Experience in Jomoke FO. *Fundamentals of Philosophy Logic and Issues in Human Existence*. Enugu: Kriscento Publishers; 2011.
24. Biodun O. Knowledge and perception of child sexual abuse in urban Nigeria: Some evidence from a community-based project. *Afr J Reprod Health*. 2010;4(2):45.
25. Terry KJ, Lalor K. Child Sexual Abuse in Sub-Saharan Africa: A Literature Review. *Child Abuse Negl*. 2004;28(1):439–460.
26. Lippert T, Cross T, Jones L, Walsh W. Telling interviewers about sexual abuse: Predictors of child disclosure at forensic interviews. *Child Maltreat*. 2009;14(1):100–113.
27. US Department of Health and Human Services. (2010). *Child maltreatment 2010* [cited 2018 Dec 10]. Available from: http://www.acf.hhs.gov/programs/cb/stats_research/index.htm#can.
28. Finkelhor D, Ormrod R, Turner H, Hamby S. School, police, and medical authority involvement with children who have experienced victimization. *Arch Pediatr Adolesc Med*. 2011;165(1):9–15.
29. Jones JS, Alexander C, Wynn BN, Rossman L, Dunnuck C. Why women don't report sexual assault to the police: the influence of psychosocial variables and traumatic injury. *J Emerg Med*. 2009;36:417–424.
30. Moher D, Liberati A, Tetzlaff J, Altman DG. The PRISMA Group Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Med*.2009;6(7):e1000097. <https://doi.org/10.1371/journal.pmed.1000097>.
31. Abdulkadir I, Musa HH, Umar LW, Musa S, Jimoh WA, Nauzo AM. Child abuse in Minna, Niger state, Nigeria. *Niger Med J*. 2011;52:79–82.
32. Daru PH, Osagie EO, Pam IC, Mutihir JT, Silas OA, Ekwempu CC. Analysis of cases of rape as seen at the Jos University Teaching Hospital, Jos, North Central Nigeria. *Niger J Clin Pract*.2011;14:47–51.
33. Ashimi AO, Amole TG1, Ugwa EA. Reported Sexual Violence among Women and Children Seen at the Gynecological Emergency Unit of a Rural Tertiary Health Facility, Northwest Nigeria. *Ann Med Health Sci Res*. 2015;5(1):26–29.
34. Bugaje MA, Ogunrinde GO, Faruk JA. Child sexual abuse in Zaria, Northwestern Nigeria. *Niger J Paediatr*. 2012;39:110–114.

35. Hassan M, Awosan KJ, Panti AA, Nasir S, Tunau K, Umar AG, et al. Prevalence and pattern of sexual assault in Usmanu Danfodiyo University Teaching Hospital, Sokoto Nigeria. *Pan Afr Med J.* 2016;24:332. doi:10.11604/pamj.2016.24.332.9462.
36. Chinawa JM, Ibekwe RC, Ibekwe MU, Obi E, Mouneke VU, Obu DC, et al. Prevalence and pattern of sexual abuse among children attending Ebonyi State University Teaching Hospital, Abakiliki, Ebonyi State. *Nig J Paed.* 2013;40(3):227–231.
37. Ekabua JE, Agan U, Ekanem EI, Itam IH, Ogaji DS. Trauma Related to Sexual Assault In Calabar, South Eastern Nigeria. *Nig J Med.* 2006;15(1):72–74.
38. Nwolisa CE, Ofondu E, Ezeogu J, Odinaka K, Adimora O. Event characteristics and socio-demographic features of rape victims presenting for post exposure prophylaxis at the Federal Medical Centre, Owerri Nigeria. *Res J Health Sci.* 2016;4(3):195–202.
39. Olatunya OS, Akintayo AA, Olofinbiyi B, Isinkaye AO, Ogundare EO, Akinboboye O. *Paediatr Int Child Health.* 2013;33(4):247–252. doi: 10.1179/2046905513Y.0000000089.
40. Akinlusi FM, Rabiou KA, Olawepo TA, Adewunmi AA, Ottun TA, Akinola OI. Sexual assault in Lagos, Nigeria: a five year retrospective review. *BMC Womens Health.* 2014;14:115. doi: 10.1186/1472-6874-14-115.
41. Badejoko OO, Anyabolu HC, Badejoko BO, Ijarotimi AO, Kuti O, Adejuyigbe EA. Sexual assault in Ile-Ife, Nigeria. *Niger Med J.* 2014;55:254–259.
42. Ige OK, Fawole OI. Evaluating the Medical Care of Child Sexual Abuse Victims in a General Hospital in Ibadan, Nigeria. *Ghana Med J.* 2012;46(1):22–26.
43. Adeleke NA, Olowookere AS, Hassan MB, Komolafe JO, Asekun-Olarinmoye EO. Sexual assault against women at Osogbo southwestern Nigeria. *Nig J Clin Pract.* 2012;15(2):190–193.
44. Ezechi OC, Musa ZA, David AN, Wapmuk AE, Gbajabiamila TA, Idigbe IE, et al. Trends and patterns of sexual assaults in Lagos south western Nigeria. *Pan Afr Med J.* 2016; 24:261. doi:10.11604/pamj.2016.24.261.9172.
45. Duru CO, Ederiane OE, Akinbami FO. Child sexual abuse: a review of cases presenting at the outpatient clinic of a tertiary health centre in Bayelsa State, Nigeria. *Global Adv Res J Med Med Sci.* 2014;3(11):354–361.
46. Akhiwu W, Umanah IN, Olueddo AN. Sexual assaults in Benin City, Nigeria. *TAF Prev Med Bull.* 2013;12(4):377–382.
47. Girgira T, Tilahun B, Bacha T. Time to presentation, pattern and immediate health effects of alleged child sexual abuse at two tertiary hospitals in Addis Ababa, Ethiopia. *BMC Public Health.* 2014;14:92. doi: 10.1186/1471-2458-14-92.
48. Ononge S, Wandabwa J, Kiondo P, Busingye R. Clinical presentation and management of alleged sexually assaulted females at Mulago hospital, Kampala, Uganda. *Afr Health Sci.* 2005; 5(1):50–54.
49. Foumane P, Dohbit JS, Monebenimp F, Natolga B, Meka ENU, Mboudou ET. Clinical Study of Rape against Females at the Yaoundé Gyneco-Obstetric and Pediatric Hospital, Cameroun. *Adv Sex Med.* 2014;4:11–16.
50. Almadani O, Bamousa M, Alsaif D, Kharoshah MAA, Alsowayigh K. Child physical and sexual abuse in Dammam, Saudi Arabia: A descriptive case-series analysis study. *Egypt J Forensic Sci.* 2012;2:33–37.
51. Mwaheb MA. Child Sexual Abuse in Fayoum Governorate, Egypt (2010-2014). *J Forensic Res.* 2016;7:332. doi: 10.4172/2157-7145.1000332.
52. Haile RT, Kebeta ND, Kassie GM. Prevalence of sexual abuse of male high school students in Addis Ababa, Ethiopia. *BMC Int Health Hum Rights.* 2013;13:24. doi: 10.1186/1472-698X-13-24.
53. Melesse F, Kassie A. Child abuse in urban setting: a one-year analysis of hospital information on abused children at Yekatit 12 hospital, Addis Ababa. *Ethiop Med J.* 2005;43(4):223–232.
54. Chesshyre E, Molyneux EM. Presentation of child sexual abuse cases to Queen Elizabeth Central Ho-

spital following the establishment of an HIV post-exposure prophylaxis programme. *Malawi Med J.* 2009;21(2):54–58.

55. Malemo KL, Lussy JP, Kimona C, Nyavandu K, Mukekulu Eugenie K, Jonathan KML, et al. Sexual violence toward children and youth in war-torn Eastern Democratic Republic of Congo. *PLoS ONE.* 2011;6:1–5.
56. Patel A, Panchal H, Piotrowski ZH, Patel D. Comprehensive medical care for victims of sexual assault: a survey of Illinois hospital emergency departments. *Contraception.* 2008;77:426–430.
57. Badejoko OO, Kuti O, Ijarotimi AO, Badejoko BO, Oke OF, Olaiya DS, et al. Have you seen a rape kit? A snapshot at the quality of care of rape survivors in Nigerian tertiary hospitals. *Trop J Obstet Gynaecol.* 2010;27(2):23–26.
58. Koss MP, Figueredo AJ, Prince RJ. Cognitive mediation of rape's mental, physical, and social health impact: Tests of four models in cross-sectional data. *J Consult Clin Psychol.* 2002; 70(4):926–941.
59. Linden JA. Care of the adult patient after sexual assault. *N Engl J Med.* 2011;365:834–841.
60. Akpoghome TU, Nwano TC. Examining the Incidences of Sexual Defilement of Children in Nigeria. *Donn J Law Conflict Resolut.* 2016;2(1):1–9.