

CASE REPORT IN INTERNAL MEDICINE AND IMMUNOLOGY

Peripheral neuropathy as initial manifestation of Churg-Strauss Syndrome

Antonio Villa¹, Marianna Gregorio², Francesca Bassi³, Ignazio Santilli³

Affiliations:

¹ *Medical Doctor, Department of Emergency, ASST Monza, Desio Hospital, Monza, Italy*

² *Medical Doctor, Department of Internal Medicine, ASST Monza, Desio Hospital, Monza, Italy*

³ *Medical Doctor, Department of Neurology, ASST Monza, Desio Hospital, Monza, Italy*

Corresponding Author:

Dr Antonio Villa, ASST Monza, PO Desio. Via Fiuggi, 56 20159 Milan, Italy. E-mail: antonio_villa@fastwebnet.it

Abstract

Peripheral neuropathy is a well-known complication of primary systemic vasculitides that are characterized by inflammatory cell infiltration causing necrosis of blood vessels. Neurological complications in eosinophilic granulomatosis with polyangiitis (EGPA), previously known as Churg-Strauss syndrome, can be seen in 60-70% of cases, and in a portion of them neurological dysfunction may precede the involvement of other organs. An early recognition of this entity is the key to successful treatment and positive outcome. We report a case of Churg-Strauss syndrome in a young man with signs of rapidly evolving peripheral neuropathy. The diagnosis was established based on the hypereosinophilia, nasal polyposis, a positive biopsy and ANCA serology, without a history of asthma. In this case report, the peripheral neuropathy was the initial manifestation of the Churg-Strauss syndrome.

KEY WORDS: Churg-Strauss syndrome; eosinophilic granulomatosis with polyangiitis; eosinophilic vasculitis; peripheral neuropathy.

Riassunto

Le neuropatie periferiche sono complicazioni ben note delle vasculiti sistemiche primarie; queste sono caratterizzate da infiltrazioni di cellule infiammatorie che causano necrosi dei vasi sanguigni. I sintomi neurologici della poliangioite con granulomatosi eosinofila, precedentemente nota come sindrome di Churg-Strauss, possono essere osservati nel 60-70% dei casi e in una parte dei casi la disfunzione neurologica può precedere il coinvolgimento di altri organi. Un riconoscimento precoce di questa entità è la chiave per ottenere un trattamento tempestivo ed esiti positivi. Segnaliamo un caso di sindrome di Churg-Strauss in un giovane con segni di neuropatia periferica in rapida evoluzione. In questo caso, la diagnosi è stata stabilita sulla base dell'ipereosinofilia, della poliposi nasale, della biopsia e della positività agli ANCA, in assenza di una storia di asma. La neuropatia periferica si è presentata come manifestazione iniziale della sindrome.

TAKE-HOME MESSAGE: Neurological involvement is frequent in Churg-Strass syndrome (CSS), and sometimes it may represent the first manifestation of this syndrome. Therefore, it is important to make early diagnosis of CSS in case of neuropathy, also in absence of the classic clinical symptoms of this syndrome.

Competing interests: none declared

Copyright © 2019 Antonio Villa et al.

Edizioni FS Publishers

This is an open access article distributed under the Creative Commons Attribution (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. See <http://www.creativecommons.org/licenses/by/4.0/>.

Cite this article as: Villa A, Gregorio M, Bassi F, Santilli I. Peripheral neuropathy as initial manifestation of Churg-Strauss Syndrome. [published online ahead of print December 15, 2019]. *J Health Soc Sci.* doi 10.19204/2019/prph5

DOI 10.19204/2019/prph5

Received: 30 Nov 2019 **Accepted:** 10 Dec 2019 **Published Online:** 15 Dec 2019

INTRODUCTION

Peripheral neuropathy is a well-known complication of primary systemic vasculitides, that are characterized by inflammatory cell infiltration causing necrosis of blood vessels. Antineutrophil cytoplasmic antibody (ANCA)-associated small vessel systemic vasculitides and medium-sized vessel vasculitis are known to frequently damage the peripheral nervous system [1-5]. In some cases, peripheral neuropathy is one of the first symptoms of systemic vasculitis [1, 2, 5-7]. Churg-Strauss syndrome (CSS) is a systemic disorder characterized by asthma, hypereosinophilia, and systemic vasculitis and frequently involves peripheral nerves and skin. Some cases of this syndrome presented without history of asthma indicating that asthma is not essential to diagnosis [8]. We describe a case of Churg-Strauss syndrome in a young man (without asthma) having signs of rapidly evolving peripheral neuropathy.

CASE REPORT

A 47-year-old man presented with a fifteen-day history of pain in his lower and upper limbs associated with paresthesias and hypostenia. His medical history revealed no history of asthma, but in the last year some episodes of obstructive-congestive rhinitis were described. His physical examination revealed 130/70 mmHg blood pressure, 74/min pulses, normal respiratory rate and oxygen saturation and no fever. The pain measured with the Numeric Pain Rating Scale (NPRS), which is a 11-point numeric scale ranging from '0' ('no pain') to '10' ('extreme pain'), was as '8'. His neurological examination showed: Left eyelid ptosis, less strenght in left deltoid and triceps, difficulty in tightening the fists bilaterally, dysesthesia with hypoesthesia from the forearm to the hand, bilaterally. The right stylus-radial, achilles and medioplant osteo-tendon reflexes were absent. There was mild deficit of proximal strength in the lower limbs. Not sensory levels. He was able to walk on points and heels.

Laboratory examination revealed white blood count 28.5×10^9 (eosinophils 61%); C-reactive protein level was 62 mg/dL (normally, 0.0-5.0 mg/dL); creatine kinase (CK) and transaminase levels were increased to 62 mg/dL and 93 mg/dL, respectively. Chest x-ray and head computed tomography scan were normal. The patient was admitted in the neurological ward. Blood exams confirmed leukocytosis with hypereosinophilia. Serum tests were positive for ANCA. Nerve conduction studies showed a neurogenic involvement of the investigated muscles, compatible with axonal motor nerve distress with blocks of motor conduction and sensory axonal impairment with asymmetric distribution; these findings were suggestive of multiplex mononeuritis. Nerve biopsy showed features of axonopathy. Muscle biopsy showed a picture with features of eosinophilic vasculitis. Spirometry was normal, with a FEV1/FVC rate of 77%

of predicted. He was treated with intravenous methyl-prednisolone and cyclophosphamide, followed by maintenance with azathioprine. After some days of therapy, a blood exam control showed: White blood count 20.5×10^9 (eosinophils 5%); C-reactive protein level 5 mg/dL; creatine kinase levels 51 mg/dL, and transaminase levels 30 mg/dL. Patient improved and there was no recurrence of symptoms during the first 2 years after the initial diagnosis.

DISCUSSION

The ANCA-associated vasculitides are a collection of relatively rare autoimmune diseases of unknown cause, characterized by inflammatory cell infiltration causing necrosis of blood vessels [9]. The association between ANCA and vasculitis was first described in 1982 [10]. The ANCA-associated vasculitides include granulomatosis with polyangiitis (GPA, formerly known as Wegener's granulomatosis), microscopic polyangiitis (MPA), and eosinophilic granulomatosis with polyangiitis (EGPA, previously known as Churg-Strauss syndrome) [9].

Named Churg-Strauss syndrome for many years, this entity has now been recognized by the 2012 revised nomenclature for vasculitides as eosinophilic granulomatosis with polyangiitis (EGPA) [11]. There are no commonly accepted diagnostic criteria for EGPA. In 1990, the American College of Rheumatology defined the classification criteria to discriminate the different vasculitides and identified six criteria for EGPA: 1. asthma; 2. eosinophilia > 10%; 3. neuropathy; 4. nonfixed lung infiltrates; 5. paranasal sinus abnormalities; 6. extravascular eosinophils on biopsy. With four or more of these criteria, vasculitis can be classified as EGPA with a sensitivity of 85% and a specificity of 99.7% [12].

The annual incidence rates of EGPA were 0.5-3.7 per million in Europe [13]. The relative rarity and non-specific presentation of the ANCA-associated vasculitides pose diagnostic challenges and often result in a significant diagnostic delay of more than 6 months in a third of patients [9].

EGPA mainly affects patients with asthma, sinusitis, allergic rhinitis, and nasal polyposis. In Table 1 the main clinical features and their prevalence are reported [14].

Table 1. Main clinical features in EGPA and their prevalence [14, modified].

| Clinical features | Prevalence (%) |
|------------------------------------|-----------------------|
| Asthma | 91-100 |
| Ear, noise, and throat involvement | 48-75 |
| Neuropathy | 55-72 |
| Pulmonary involvement | 65-91 |
| Cutaneous involvement | 40-52 |
| Renal involvement | 27 |
| Cardiac involvement | 27-35 |
| Gastrointestinal involvement | 23-32 |
| Central nervous system involvement | 5-9 |
| ANCA positivity | 38 |

Neurological symptoms can be seen in 60-70% of patients with EGPA, secondary only to pulmonary manifestations in the overall frequency of organ involvement [15]. Some of the neurological manifestations in EGPA include cranial neuropathy, extremity mononeuropathy, multiplex mononeuropathy, symmetrical length-dependent polyneuropathy, and rarely manifestation of central nervous system dysfunction [16, 17]. In a portion of cases, neurological dysfunction may precede the involvement of other organs or be part of the presenting

constellation [5]. An early recognition of this entity is key to leading to timely treatment and positive outcome [17]. Churg and Strauss reported three primary histopathological alterations: Eosinophilic tissue infiltration, necrotizing vasculitis and extravascular granulomas. The inflammatory process in Churg-Strauss syndrome tends to affect smaller epineural arterioles than those in systemic vasculitis [18]. The underlying pathogenic mechanisms of vasculitic neuropathy are still not fully understood. Presumably, the interplay of humoral antibodies (ANCA), activated neutrophil granulocytes, the complement system, and endothelial cells could lead to inflammation and luminal narrowing of the vessel wall of small epineural arteries, resulting in ischemic lesions of the peripheral nerve [5, 19]. Therefore, the neuropathy is caused mainly by nerve ischemia due to occlusion of *vasa nervorum*. The result of this infarction is a loss of sensory and motor axons [18].

EGPA is traditionally described to evolve through a prodromic, allergic phase characterized by asthma and rhinosinusitis, an eosinophilic phase hallmarked by peripheral eosinophilia and organ involvement, and a vasculitic phase with clinical manifestations due to small-vessel vasculitis [20]. These phases partially overlap and may not appear in such a defined order, although asthma and rhinosinusitis only rarely arise after the vasculitic manifestations [21, 22]. Asthma is found in 95-100% of patients and may precede the systemic disease manifestations of many years [20].

The vasculitic phase is heralded by constitutional symptoms (e.g., fever, weight loss, fatigue) and often by an apparently paradoxical improvement of asthma. Peripheral neuropathy is a cardinal feature of this phase, affecting 70% of the patients [2, 20, 23]. It is characterized by axonal damage on electrophysiological studies and frequently affects the peroneal, tibial, ulnar, and

median nerves; the most common pattern is multiplex mononeuritis; sensory deficits and neuropathic pain are also frequent [2, 20]. The diagnosis of polyneuropathy is based on clinical and electrophysiologic studies, but precise histology, immunohistochemistry and morphometric studies of the peripheral nerve biopsy may be decisive to make diagnosis [18, 24]. Nerve recovery is very slow, and only partial, thus early treatment is essential in preventing extensive damage and reduced quality of life [25].

In rare cases, vasculitis is confined to peripheral nerves without evidence of vasculitis in other organ systems [5]. In this case, the diagnosis of EGPA was established based on the hypereosinophilia, nasal polyposis, a positive biopsy and ANCA serology. This case indicates that the hypereosinophilia is the main risk factor for EGPA without the history of asthma. Several investigators reported that EGPA might occur in patients without asthma [8, 26-32]. Therefore, the absence of asthma should be not an absolute criterium to exclude the diagnosis of EGPA. These cases, which did not fulfill the criteria of classic EGPA, were also termed variant forms or incomplete forms ('formes frustes') of EGPA [30]. Although Churg-Strauss syndrome is rare, clinicians can ill afford to misdiagnose it, since it is easily diagnosable and treatable and potentially fatal in the absence of treatment.

Identification of cause is one of the important goals of clinician's approach to a patient with newly diagnosed peripheral neuropathy. Evaluating a patient with neuropathy by thorough history taking, clinical examination, and relevant laboratory investigations for presence of any associated medical conditions, which may serve as a clue to the diagnosis, is an important tool to accomplish this goal [31]. A structured approach based on a careful clinical and electrodiagnostic

assessment with attention to the pattern of nerve involvement can help narrow the differential diagnosis and rationalise laboratory evaluation [31]. Churg-Strauss syndrome frequently presents polyneuropathy as a complication; since its remission depends on immunosuppression therapy, it is important to recognize it at an early stage.

References

1. de Groot K, Schmidt DK, Arlt AC, Gross WL, Reinhold-Keller E. Standardized neurologic evaluations of 128 patients with Wegener's granulomatosis. *Arch Neurol.* 2001;58(8):1215-1221.
2. Cattaneo L, Chierici E, Pavone L, Grasselli C, Manganeli P, Buzio, et al. Peripheral neuropathy in Wegener's granulomatosis, Churg-Strauss syndrome and microscopic polyangiitis. *J Neurol Neurosurg Psychiatry.* 2007;78(10):1119-1123.
3. Pagnoux C, Seror R, Henegar C, Mahr R, Cohen P, Le Guern V, et al. Clinical features and outcomes in 348 patients with polyarteritis nodosa: a systematic retrospective study of patients diagnosed between 1963 and 2005 and entered into the French Vasculitis Study Group Database. *Arthritis Rheum.* 2010;62(2):616-626. doi:10.1002/art.27240.
4. Mahr A, Katsahian S, Varet H, Guillenin L, Hagen EL, Höglund P, et al. Revisiting the classification of clinical phenotypes of anti-neutrophil cytoplasmic antibody-associated vasculitis: a cluster analysis. *Ann Rheum Dis.* 2013;72(6):1003-1010. doi: 10.1136/annrheumdis-2012-201750.

5. Wolf J, Schmitt V, Palm F, Gran AJ, Bergner R. Peripheral neuropathy as initial manifestation of primary systemic vasculitides. *J Neurol*. 2013;260(4):1061-1070. doi: 10.1007/s00415-012-6760-7.
6. Zhang W, Zhou G, Shi Q, Zhang X, Zeng XF, Zhang FC. Clinical analysis of nervous system involvement in ANCA-associated systemic vasculitides. *Clin Exp Rheumatol*. 2009;27(1 Suppl):S65-69.
7. Suppiah R, Hadden RD, Batra R, Arden NK, Collins MP, Guillenin L, et al. Peripheral neuropathy in ANCA-associated vasculitis: outcomes from the European Vasculitis Study Group trials. *Rheumatology (Oxford)*. 2011;50(12):2214-2222. doi: 10.1093/rheumatology/ker266.
8. Chen KR, Ohata Y, Sakurai M, Nakayama H. Churg-Strauss syndrome: report of a case without preexisting asthma. *J Dermatol*. 1992;19(1):40-47.
9. Yates M, Watts R. ANCA-associated vasculitis. *Clin Med (Lond)*. 2017;17(1):60-64. doi: 10.7861/clinmedicine.17-1-60.
10. Davies DJ, Moran JE, Niall JF, Ryan GB. Segmental necrotising glomerulonephritis with antineutrophil antibody: possible arbovirus aetiology? *Br Med J*. 1982;285(6342):606. doi: <https://doi.org/10.1136/bmj.285.6342.606>.
11. Jennette JC, Falk RJ, Bacon PA, Basu N, Cid MC, Ferrario F, et al. 2012 revised international Chapel Hill Consensus Conference Nomenclature of Vasculitides. *Arthritis Rheum*. 2013;65(1):1-11. doi: 10.1002/art.37715.

12. Masi AT, Hunder GG, Lie JT, Michel BA, Bloch DA, Arend WP, et al. The American College of Rheumatology 1990 criteria for the classification of Churg-Strauss syndrome (allergic granulomatosis and angiitis). *Arthritis Rheum.* 1990;33(8):1094-1100.
13. Watts RA, Mahr A, Mohammad AJ, Gatenby P, Basu N, Flores-Suarez LF. Classification, epidemiology and clinical subgrouping of antineutrophil cytoplasmic antibody (ANCA)-associated vasculitis. *Nephrol Dial Transplant.* 2015;30(Suppl 1):i14-22. doi: 10.1093/ndt/gfv022.
14. Gioffredi A, Maritati F, Oliva E, Buzio C. Eosinophilic granulomatosis with polyangiitis: an overview. *Front Immunol.* 2014;5:549.
15. Gwathmey KG, Burns TM, Collins MP, Dick JB. Vasculitic neuropathies. *Lancet Neurol.* 2014;13(1):67-82. doi: [https://doi.org/10.1016/S1474-4422\(13\)70236-9](https://doi.org/10.1016/S1474-4422(13)70236-9).
16. Wolf J, Bergner R, Mutallib S, Buggle F, Grau AJ. Neurologic complications of Churg-Strauss syndrome – a prospective monocentric study. *Eur J Neurol.* 2010;17(4):582-588. doi: 10.1111/j.1468-1331.2009.02902.x.
17. Santos-Pinheiro F, Li Y. Eosinophilic granulomatosis with polyangiitis (Churg-Strauss syndrome) presenting with polyneuropathy - a case series. *J Clin Neuromusc Dis.* 2015;16(3):125-130. doi: 10.1097/CND.0000000000000068.
18. Bougea A, Anagnostou E, Spandideas N, Triantafylou N, Karakizou E. An update of neurological manifestations of vasculitides and connective tissue diseases: a literature

- review. *Einstein (Sao Paulo)*. 2015;13(4):627-635. doi: 10.1590/s1679-45082015RW3308.
19. Kallenberg CG. Pathophysiology of ANCA-associated small vessel vasculitis. *Curr Rheumatol Rep*. 2010;12(6):399-405. doi: 10.1007/s11926-010-0138-6.
20. Vaglio A, Buzio C, Zwerina J. Eosinophilic granulomatosis with polyangiitis (Churg-Strauss): state of the art. *Allergy*. 2013;68(3):261-273. doi: 10.1111/all.12088.
21. Endo T, Katsuta Y, Kimura Y, Kikuchi A, Aramaki T, Takano T, et al. A variant form of Churg-Strauss syndrome: initial temporal non-giant cell arteritis followed by asthma – is this a distinct clinicopathologic entity? *Hum Pathol*. 2000;31(9):1169-1171.
22. Noth I, Streck ME, Leff AR. Churg-Strauss syndrome. *Lancet*. 2003;361(9357):587-594.
23. Sinico RA, Di Toma L, Maggiore U, Bottero P, Radice A, Tosoni C, et al. Prevalence and clinical significance of antineutrophil cytoplasmic antibodies in Churg-Strauss syndrome. *Arthritis Rheum*. 2005;52(9):2926-2935.
24. Sablé-Fourtassou R, Cohen P, Mahr A, Pagnoux C, Mouthon L, Jayne D, et al. Antineutrophil cytoplasmic antibodies and the Churg-Strauss syndrome. *Ann Intern Med*. 2005;143(9):632-638.
25. Rutgers A, Kallenberg CG. Vasculitis syndromes: peripheral neuropathy in AAV – when vasculitis hits a nerve. *Nat Rev Rheumatol*. 2011;8(3):127-128. doi: 10.1038/nrrheum.2011.184.

26. Sasaki A, Hasegawa M, Nakazato Y, Ishida Y, Saltoh S. Allergic granulomatosis and angiitis (Churg-Strauss syndrome). Report of an autopsy case in a nonasthmatic patient. *Acta Pathol Jpn.* 1988;38(6):761-768.
27. Churg A. Recent advances in the diagnosis of Churg-Strauss syndrome. *Mod Pathol.* 2001;14(12):1284-1293.
28. Malik TQ, Youmbissi TJ, Gacha R, Abdelraham M, Al-Khursany AL, Karkar A. Atypical presentation of Churg-Strauss syndrome: another “forme fruste” of the disease? *Am J Med Sci.* 2002;324(5):276-278.
29. Sevinc A, Hasanoglu HC, Gokirmak M, Yldirim Z, Baysal T, Mizrak B. Allergic granulomatous and angiitis in the absence of asthma and blood eosinophilia: a rare presentation of limited Churg-Strauss syndrome. *Rheumatol Int.* 2004;24(5):301-304.
30. Kajihara H, Tachiyama Y, Hirose T, Takada A, Saito K, Murai T, et al. Eosinophilic coronary periarteritis (vasospastic angina and sudden death), a new type of coronary arteritis: report of seven autopsy cases and a review of the literature. *Virchows Arc.* 2013;462(2):239-248. doi: 10.1007/s00428-012-1351-7.
31. Pandita KK, Bhat KJ, Razdan S, Kudyar RP. “Peripheral neuropathy crippling bronchial asthma”: two rare case reports of Churg-Strauss syndrome. *Case Rep Pulmonol.* 2014;2014:673906. doi:10.1155/2014/673906.

32. Zeng M, Liu X, Liu Y. Eosinophilic granulomatosis with polyangiitis presenting with skin rashes, eosinophilic cholecystitis, and retinal vasculitis. *Am J Case Rep.* 2016;17:864-868.